









---

A MONTE CARLO STUDY OF ROBUST LOCATION  
ESTIMATES WITH NON-GAUSSIAN MULTIVARIATE DATA

by

M. Anthony Wong and Gregory M. Lenhart

Working Paper #1135

August 1980

---



A MONTE CARLO STUDY OF ROBUST LOCATION  
ESTIMATES WITH NON-GAUSSIAN MULTIVARIATE DATA

by

M. Anthony Wong and Gregory M. Lenhart

Working Paper #1135

August 1980





A MONTE CARLO STUDY OF ROBUST LOCATION ESTIMATES  
WITH NON-GAUSSIAN MULTIVARIATE DATA

M. ANTHONY WONG and GREGORY M. LENHART  
Sloan School of Management, Massachusetts Institute of Technology,  
Cambridge, MA 02139

Four robust estimators of multivariate locations are compared in a Monte Carlo study by examining their empirical efficiency performance on a range of bivariate distributions with heavier tails than the Gaussian distribution. Using three different multivariate measures of relative efficiency, the simulation results show that the Hodges-Lehmann estimator is "safest" among the robust estimators, in that it has uniformly high efficiencies across distributions ranging from the uniform distribution on a square to the bivariate Cauchy distribution.

KEYWORDS: Monte Carlo study, robust estimators, multivariate location, multivariate relative efficiency.



## 1. INTRODUCTION

In robust estimation of location, the main concern is to obtain estimates of location which are insensitive to outliers. For the univariate case, problems and methods of robust estimation have received considerable attention (see, for example, Tukey (1960); Hodges and Lehmann (1963); Huber (1964, 1972); Bickel (1965); Andrews et al (1972)). Various estimates have been proposed and their relative behaviors have been studied for a range of symmetrical distributions with increasingly heavier tails than the Gaussian distribution.

The problem of robust estimation of multivariate location has received some attention in the literature (see Mood (1941); Bickel (1964); Gentleman (1965)). The usual estimate of multivariate location is the sample mean vector  $\bar{x}$ , whose elements are just the univariate means, and except for the estimator proposed by Gentleman (1965) the multivariate robust estimators are just vectors of univariate robust estimators obtained by analyzing the observations on each of the variables separately. Gnanadesikan and Kettenring (1972) give a survey of existing robust estimators of multivariate location and present the results of a Monte Carlo study of the following estimators:

- (i)  $x_M$ , the vector of medians of the observations on each variable, as suggested by Mood (1941);
- (ii)  $x_{HL}$ , the vector of Hodges-Lehmann estimators (i.e., the median of averages of pairs of observations, including self-pairs) for each variable, as proposed and studied by Bickel (1964);
- (iii)  $x_{T(\alpha)}$ , the vector of  $\alpha$ -trimmed means (i.e., the mean of the data remaining after omitting a proportion  $\alpha$  of the smallest and of the largest observations) for each variable.

However, their study was limited to the Gaussian case and hence the relative behaviors of the estimators for other alternative distributions have not been studied. Here the scope of their study of robust estimators of multivariate



location is extended to the non-Gaussian cases, so that the estimators can be compared across a range of distributions with thicker tails than the Gaussian distribution. The simulation experiments in this Monte Carlo study are described in Section 2, while the results are given in Section 3. Using three different multivariate measures of relative efficiencies (which differ in their sensitivities to the underlying correlation structure), we conclude that the Hodges-Lehmann estimator is preferred to the other estimators, especially when the variates are not highly correlated; a result which is in agreement with the theoretical considerations of Bickel (1964, 1965).

## 2. THE MONTE CARLO STUDY

In addition to the usual sample mean vector  $\bar{x}$ , we consider the robust alternatives: (i) the vector of medians  $x_M$ , (ii) the vector of Hodges-Lehmann estimates  $x_{HL}$ , (iii) the vector of 25% - trimmed means  $x_{T(.25)}$ , and (iv) the vector of 10% - trimmed means  $x_{T(.10)}$  as estimates of multivariate location. The performance of these five estimates were compared using generated bivariate data drawn from a range of distributions. The null sampling distribution is the bivariate Gaussian distribution  $(N(\underline{0}, \Gamma), \Gamma = \begin{bmatrix} 1 & \rho \\ \rho & 1 \end{bmatrix})$ . Other distributions considered are those with thicker tails than the Gaussian (Laplace, contaminated Gaussian  $(0.9N(\underline{0}, \Gamma) + 0.1N(\underline{0}, 9\Gamma))$ , and Cauchy) and one shorter-tailed than the Gaussian ( $\text{uniform}^2 [-1/2, 1/2]$ ). Simulated samples from these distributions can be obtained as done by Andrews et al. (1972) for the univariate case.

Three sample sizes ( $n = 20, 40$ , and  $60$ ) and three different correlations ( $\rho = 0.0, 0.5$ , and  $0.9$ ) are used in the empirical comparisons of the proposed location estimators. One hundred simulated bivariate samples are produced for each case; independent samples are drawn for  $n = 20, 40$ , and  $60$  for  $\rho = 0.0$ , and then transformed by replacing each value of the second variate



with  $\rho x_{1i} + (1 - \rho^2)^{1/2} x_{2i}$ ;  $i = 1, 2, \dots, n$  to obtain the data for  $\rho = 0.5$  and  $0.9$ . Hence, it should be noted that although the results for different sample sizes are independent, the comparisons for fixed  $n$  are not.

For each estimator (mean, median, Hodges-Lehmann, and trimmed means [ $\alpha = 0.25$  and  $0.10$ ]), the Monte Carlo mean vector and elements of the Monte Carlo covariance matrix (var (1), var (2), cov (1,2), corr (1,2), and the characteristic roots  $C_{\min}$  and  $C_{\max}$ ) are computed. Also computed are three multivariate measures of relative efficiencies of the robust estimators with respect to  $\bar{x}$ ; the measures of efficiency are (i) the square root of the product, (ii) the sum, and (iii) the square root of the sum of squares, of the characteristic roots of the Monte Carlo covariance matrix of the location estimators, and the relative efficiencies DET, TR, and TR2 are then respectively the corresponding ratios of the values for  $\bar{x}$  to the values for the various robust estimators. While DET is sensitive to the correlation  $\rho$  (decreases when  $\rho$  increases, in general), TR is independent of  $\rho$  and TR2 depends on  $\rho$  but not as sensitively as DET (especially when  $|\rho| \rightarrow 1$ ). The DET measure was proposed by Bickel (1964) and the other two are suggested by Gnanadesikan and Kettenring (1972).

### 3. RESULTS

For  $n=60$ , the Monte Carlo mean vectors of the location estimators for the symmetrical distributions are given in Table I. As expected, the entries in this table indicate that the location estimators are all reasonably unbiased (except for the case when the mean vector is used to estimate the location of the Cauchy distribution); the results for  $n=20$  and  $40$  give similar indication and are not included here.

In Table II through VI, the Monte Carlo efficiencies of the estimators when  $n=60$  are given for the Gaussian, Laplace, contaminated Gaussian, Cauchy





and uniform distributions respectively. In the Gaussian case (Table II), it is clear that there are considerable differences among the efficiencies, except for  $X_{HL}$  and  $X_{T(.10)}$  which behave much alike while consistently outperforming the other two robust alternatives (with the median vector  $X_M$  being least efficient). For correlated Gaussian data, note that the correlations between the elements of the robust estimators (especially the median  $X_M$ ) are in general less than the correlations for the sample mean vector; and this is also evident in all the non-Gaussian cases.

With Laplace data (Table III), all the robust estimates perform better than the mean vector. The median  $X_M$ , by virtue of its properties as the maximum likelihood estimator, gives the best performance here. The 25% - trimmed mean  $X_{T(.25)}$  is next best, while the Hodges-Lehman estimate  $X_{HL}$  consistently outperforms the 10% - trimmed mean  $X_{T(.10)}$ . When Gaussian data are contaminated by 10% outliers, again all robust estimates outperform the mean (Table IV); however, when the value of  $\rho$  is high, the median does not appear to be superior than the mean  $\bar{x}$ . The two estimators  $X_{HL}$  and  $X_{T(.10)}$  again behave similarly, and again give the best efficiencies among the robust estimators. For the extremely heavy-tailed Cauchy data (Table V), the median  $X_M$  and the 25% - trimmed mean  $X_{T(.25)}$  are most efficient, and  $X_{HL}$  is notably better than  $X_{T(.10)}$ . In the short-tailed uniform case (Table VI), the mean is naturally the most efficient estimator; but the Hodges-Lehmann estimator is not much worse. Thus the Hodges-Lehman estimator  $X_{HL}$  appears to have good efficiency properties for a range of symmetrical distributions (even when the distribution is short-tailed), and this is true for all the three sample sizes  $n=20, 40$  and  $60$  considered although the results for  $n=20$  and  $40$  are not shown.

To facilitate further comparisons among the robust alternatives, the three multivariate measures of relative efficiency DET, TR, and TR2 are computed



for each distribution (Tables VII through XI respectively for the Gaussian, Laplace, contaminated Gaussian, Cauchy and Uniform distributions). The results given in these tables are consistent with the observations made earlier. The median  $X_M$  is good in special cases (see Table VIII (Laplace) and Table X (Cauchy)); but it is the least efficient estimator in the Gaussian case (less than 70% for the DET measure), especially when  $\rho = 0.9$ . The Hodges-Lehmann estimator  $X_{HL}$  and the 10% - trimmed mean perform similarly; they have high efficiencies in the Gaussian case (generally greater than 85% for all three measures of relative efficiency: see Table VII) and in the uniform case, while being very resistant to outliers in the heavy-tail cases.

As pointed out earlier, the DET measure is very sensitive to the underlying correlation structure  $\rho$ . It can be shown that DET is directly proportional to  $(1-\rho^2_{\text{mean}})/(1-\rho^2_{\text{robust}})$ , where  $\rho_{\text{mean}}$  is the correlation between the elements of the sample mean vector and  $\rho_{\text{robust}}$  is the corresponding correlation for a robust estimator. Since the components of the vector of robust estimates (especially the median  $X_M$ ) are not as highly correlated as the sample mean vector (especially when the underlying correlation  $|\rho|$  is high), a higher  $|\rho|$  value will mean a smaller value of DET for a robust estimator. In fact, Bickel (1964) pointed out that when  $|\rho| \rightarrow 1$ , there exist extreme cases where even the Hodges-Lehmann estimator might have arbitrarily small relative efficiency DET. Due to the limited scope of this study, no such extreme cases are given here. However, it seems fair to conclude that the Hodges-Lehmann estimator have uniformly high efficiencies for a wide range of distribution, even when the correlation  $\rho$  is as high as 0.9. Thus the simulated results are in agreement with a straightforward extension of the univariate asymptotics obtained by Bickel (1965) for the Hodges-Lehmann estimator.

#### 4. DISCUSSION

In this paper, the results of a Monte Carlo study of the robust estimators



considered by Gnanadesikan and Kettenring (1972) are presented. The estimators are compared using three different multivariate measures of relative efficiency, across a range of bivariate distributions with heavier tails than the Gaussian distribution. The results indicate that the Hodges-Lehmann estimator proposed by Bickel (1964) is the "safest" among the robust alternatives; a result which is in agreement with Bickel's (1964, 1965) asymptotic considerations. Thus, we conclude that the Hodges-Lehmann estimate is to be preferred (when computationally feasible -- it requires the largest number of calculations among the robust alternatives) where the amount of contamination and the distributional form is not known precisely. However, it should be pointed out that there exist extreme cases (when  $|\rho|$  is very close to 1) where this estimate behaves very poorly (see Bickel, 1964).



TABLE I  
MONTE CARLO MEANS OF LOCATION ESTIMATORS\*  
(n=60)

Sampling Distribution		$\bar{x}$	$x_M$	$x_{HL}$	$x_{T(.25)}$	$x_{T(.10)}$
Gaussian	$\rho=0.0$	-0.0140	-0.0303	-0.0142	-0.0192	-0.0121
		0.0205	0.0117	0.0180	0.0140	0.0175
	$\rho=0.5$	0.0107	0.0008	0.0069	0.0052	0.0069
	$\rho=0.9$	-0.0036	-0.0161	-0.0051	-0.0072	-0.0038
Laplace	$\rho=0.0$	-0.0207	-0.0199	-0.0136	-0.0165	-0.0107
		0.0339	0.0127	0.0202	0.0125	0.0254
	$\rho=0.5$	0.0190	-0.0020	0.0078	0.0039	0.0106
	$\rho=0.9$	-0.0038	-0.0139	-0.0060	-0.0094	-0.0029
Contaminated Gaussian	$\rho=0.0$	-0.0163	-0.0319	-0.0163	-0.0198	-0.0142
		0.0161	0.0151	0.0220	0.0176	0.0214
	$\rho=0.5$	0.0058	-0.0003	0.0076	0.0059	0.0079
	$\rho=0.9$	-0.0077	-0.0162	-0.0070	-0.0072	-0.0038
Cauchy	$\rho=0.0$	-1.2784	-0.0253	-0.0207	-0.0180	-0.0171
		-1.7073	0.0181	0.0320	0.0216	0.0495
	$\rho=0.5$	-2.1178	-0.0027	-0.0065	-0.0037	-0.0151
	$\rho=0.9$	-1.8948	-0.0198	-0.0233	-0.0184	-0.0175
Uniform	$\rho=0.0$	0.0028	0.0053	0.0030	0.0032	0.0033
		-0.0049	-0.0058	-0.0059	-0.0082	-0.0067
	$\rho=0.5$	-0.0028	-0.0061	-0.0035	-0.0041	-0.0034
	$\rho=0.9$	0.0004	0.0033	0.0003	0.0001	0.0003

\*The first elements of the estimates do not vary with  $\rho$  and therefore are included only for the case  $\rho=0$ .





TABLE II

MONTE CARLO EFFICIENCIES OF LOCATION ESTIMATES\*  
(Gaussian Distribution;  $n=60$ )

	$\bar{x}$	$X_M$	$X_{HL}$	$X_T(.25)$	$X_T(.10)$
$\rho=0.0$					
var(1)	0.0137	0.0245	0.0150	0.0184	0.0149
var(2)	0.0148	0.0222	0.0156	0.0170	0.0160
cov(1,2)	-0.0030	-0.0027	0.0029	-0.0026	0.0025
corr(1,2)	-0.2106	-0.1164	-0.1887	-0.1493	-0.1636
$C_{min}$	0.0112	0.0204	0.0124	0.0150	0.0129
$C_{max}$	0.0172	0.0263	0.0182	0.0205	0.0181
$\rho=0.5$					
var(2)	0.0119	0.0179	0.0129	0.0145	0.0132
cov(1,2)	0.0042	0.0050	0.0045	0.0052	0.0045
corr(1,2)	0.3327	0.2384	0.3243	0.3207	0.3200
$C_{min}$	0.0084	0.0152	0.0093	0.0109	0.0095
$C_{max}$	0.0171	0.0272	0.0186	0.0221	0.0186
$\rho=0.9$					
var(2)	0.0115	0.0232	0.0123	0.0156	0.0124
cov(1,2)	0.0110	0.0166	0.0118	0.0144	0.0118
corr(1,2)	0.8760	0.6942	0.8660	0.8495	0.8670
$C_{min}$	0.0015	0.0073	0.0018	0.0025	0.0018
$C_{max}$	0.0236	0.0405	0.0255	0.0315	0.0255

\*var(1) does not change with  $\rho$ .



TABLE III

MONTE CARLO EFFICIENCIES OF LOCATION ESTIMATES\*  
(Laplace Distribution; n=60)

	$\bar{x}$	$X_M$	$X_{HL}$	$X_{T(.25)}$	$X_{T(.10)}$
$\rho=0.0$					
var(1)	0.0334	0.0223	0.0237	0.0228	0.0269
var(2)	0.0270	0.0174	0.0193	0.0178	0.0215
cov(1,2)	-0.0070	-0.0028	-0.0046	-0.0044	-0.0055
corr(1,2)	-0.2346	-0.1436	-0.2130	-0.2169	-0.2298
C <sub>min</sub>	0.0225	0.0161	0.0164	0.0153	0.0181
C <sub>max</sub>	0.0379	0.0236	0.0266	0.0253	0.0304
$\rho=0.5$					
var(2)	0.0225	0.0137	0.0170	0.0160	0.0186
cov(1,2)	0.0106	0.0048	0.0068	0.0052	0.0074
corr(1,2)	0.3867	0.2727	0.3385	0.2737	0.3302
C <sub>min</sub>	0.0160	0.0116	0.0128	0.0132	0.0143
C <sub>max</sub>	0.0399	0.0244	0.0279	0.0256	0.0313
$\rho=0.9$					
var(2)	0.0267	0.0191	0.0178	0.0165	0.0205
cov(1,2)	0.0270	0.0136	0.0181	0.0162	0.0209
corr(1,2)	0.9045	0.6588	0.8811	0.8340	0.8888
C <sub>min</sub>	0.0028	0.0070	0.0024	0.0032	0.0026
C <sub>max</sub>	0.0572	0.0343	0.0391	0.0361	0.0449

\*var(1) does not change with  $\rho$ .



TABLE IV

MONTE CARLO EFFICIENCIES OF LOCATION ESTIMATES\*  
(Contaminated Gaussian Distribution; n=60)

	$\bar{x}$	$x_M$	$x_{HL}$	$x_{T(.25)}$	$x_{T(.10)}$
$\rho = 0.0$					
var(1)	0.0259	0.0288	0.0196	0.0219	0.0189
var(2)	0.0273	0.0261	0.0191	0.0195	0.0194
cov(1,2)	-0.0044	-0.0030	-0.0032	-0.0026	-0.0029
corr(1,2)	-0.1658	-0.1080	-0.1666	-0.1259	-0.1519
$C_{min}$	0.0221	0.0242	0.0161	0.0178	0.0162
$C_{max}$	0.0311	0.0307	0.0226	0.0235	0.0221
$\rho = 0.5$					
var(2)	0.0231	0.0199	0.0165	0.0176	0.0167
cov(1,2)	0.0091	0.0059	0.0053	0.0060	0.0054
corr(1,2)	0.3730	0.2471	0.2915	0.3054	0.3022
$C_{min}$	0.0153	0.0170	0.0126	0.0134	0.0123
$C_{max}$	0.0337	0.0317	0.0236	0.0261	0.0233
$\rho = 0.9$					
var(2)	0.0227	0.0275	0.0162	0.0185	0.0156
cov(1,2)	0.0214	0.0198	0.0153	0.0170	0.0149
corr(1,2)	0.8819	0.7033	0.8593	0.8467	0.8692
$C_{min}$	0.0029	0.0083	0.0025	0.0031	0.0022
$C_{max}$	0.0457	0.0479	0.0333	0.0373	0.0323

\*var(1) does not change with  $\rho$ .



TABLE V  
MONTE CARLO EFFICIENCIES OF LOCATION ESTIMATES\*  
(Cauchy Distribution; n=60)

	$\bar{x}$	$X_M$	$X_{HL}$	$X_T(.25)$	$X_T(.10)$
$\rho=0.0$					
var(1)	217.5	0.0443	0.0675	0.0496	0.1067
var(2)	378.0	0.0361	0.0477	0.0382	0.0724
cov(1,2)	247.9	-0.0064	-0.0139	0.0096	-0.0186
corr(1,2)	0.864	-0.1588	-0.2463	-0.2210	-0.2118
$C_{min}$	37.16	0.0326	0.0405	0.0327	0.0642
$C_{max}$	558.4	0.0478	0.0747	0.0551	0.1149
$\rho=0.5$					
var(2)	552.6	0.0310	0.0425	0.0350	0.0668
cov(1,2)	323.5	0.0096	0.0175	0.0123	0.0292
corr(1,2)	0.933	0.2601	0.3271	0.2948	0.3455
$C_{min}$	20.76	0.0259	0.0335	0.0280	0.0514
$C_{max}$	749.4	0.0494	0.0765	0.0566	0.1221
$\rho=0.9$					
var(2)	442.5	0.0407	0.0477	0.0372	0.0758
cov(1,2)	303.8	0.0298	0.0493	0.0358	0.0795
corr(1,2)	0.979	0.7016	0.8691	0.8343	0.8837
$C_{min}$	6.028	0.0127	0.0073	0.0070	0.0103
$C_{max}$	645.0	0.7239	0.1079	0.0798	0.1723

\*var(1) does not change with  $\rho$ .





TABLE VI

MONTE CARLO EFFICIENCIES OF LOCATION ESTIMATES  
(Uniform Distribution; n=60)

	$\bar{x}$	$X_M$	$X_{HL}$	$X_{T(.25)}$	$X_{T(.10)}$
$\rho=0.0$					
var(1)	0.0013	0.0036	0.0015	0.0025	0.0018
var(2)	0.0017	0.0047	0.0019	0.0033	0.0023
cov(1,2)	-0.0003	-0.0005	-0.0003	-0.0006	-0.0004
corr(1,2)	-0.1987	-0.1325	-0.1898	-0.1968	-0.1966
$C_{min}$	0.0011	0.0034	0.0013	0.0022	0.0016
$C_{max}$	0.0018	0.0049	0.0021	0.0036	0.0026
$\rho=0.5$					
var(2)	0.0013	0.0030	0.0016	0.0021	0.0016
cov(1,2)	0.0004	0.0005	0.0004	0.0004	0.0005
corr(1,2)	0.3026	0.1460	0.2779	0.1922	0.2785
$C_{min}$	0.0009	0.0027	0.0011	0.0018	0.0012
$C_{max}$	0.0017	0.0038	0.0020	0.0028	0.0022
$\rho=0.9$					
var(2)	0.0011	0.0026	0.0014	0.0019	0.0015
cov(1,2)	0.0010	0.0025	0.0012	0.0019	0.0014
corr(1,2)	0.8563	0.8040	0.8625	0.8622	0.8658
$C_{min}$	0.0002	0.0006	0.0002	0.0003	0.0002
$C_{max}$	0.0023	0.0056	0.0027	0.0042	0.0031

\*var(1) does not change with  $\rho$ .



TABLE VII

MONTE CARLO RELATIVE EFFICIENCIES  
(Gaussian Distribution)

(a) n=20	$X_M$	$X_{HL}$	$X_{T(.25)}$	$X_{T(.10)}$
RHO=0.0				
DET	.689909	.929654	.824216	.945078
TR	.823092	.962997	.904724	.970725
TR2	.665828	.925114	.813012	.939589
RHO=0.5				
DET	.675513	.918211	.818918	.947645
TR	.854516	.967430	.923187	.976439
TR2	.771218	.947330	.874919	.957044
RHO=0.9				
DET	.489929	.878544	.766510	.909420
TR	.866173	.978745	.941022	.979451
TR2	.817547	.968741	.903438	.965734
(b) n=40	$X_M$	$X_{HL}$	$X_{T(.25)}$	$X_{T(.10)}$
RHO=0.0				
DET	.642301	.920596	.825578	.913418
TR	.800480	.956498	.901624	.950937
TR2	.639352	.909672	.801663	.896022
RHO=0.5				
DET	.569563	.895414	.755092	.887636
TR	.773948	.950629	.874923	.946655
TR2	.624887	.910192	.773746	.902845
RHO=0.9				
DET	.387351	.821974	.635491	.797927
TR	.742013	.931933	.844873	.922730
TR2	.589295	.874864	.725732	.858875
(c) n=60	$X_M$	$X_{HL}$	$X_{T(.25)}$	$X_{T(.10)}$
RHO=0.0				
DET	.598836	.923769	.791889	.910991
TR	.779802	.963593	.894963	.958818
TR2	.616922	.932903	.809511	.927146
RHO=0.5				
DET	.591218	.913816	.776972	.904815
TR	.776325	.957170	.881217	.953769
TR2	.612270	.918062	.776205	.913590
RHO=0.9				
DET	.352009	.889731	.676133	.894352
TR	.726009	.959973	.860139	.961085
TR2	.575821	.925957	.749552	.927729



TABLE VIII  
MONTE CARLO RELATIVE EFFICIENCIES  
(Laplace Distribution)

(a) n=20	$X_M$	$X_{HL}$	$X_{T(.25)}$	$X_{T(.10)}$
RHO=0.0				
DET	1.85396	1.45846	1.63008	1.32251
TR	1.35118	1.20997	1.28091	1.15086
IR2	1.79951	1.46945	1.65121	1.32638
RHO=0.5				
DET	1.51962	1.43344	1.49287	1.30922
TR	1.25543	1.19607	1.23408	1.14116
IR2	1.61594	1.42873	1.54326	1.29782
RHO=0.9				
DET	1.14343	1.24008	1.31397	1.24067
TR	1.32428	1.17222	1.22177	1.13647
IR2	1.93295	1.39505	1.52192	1.29868
(b) n=40	$X_M$	$X_{HL}$	$X_{T(.25)}$	$X_{T(.10)}$
RHO=0.0				
DET	1.68281	1.46666	1.61039	1.31783
TR	1.27819	1.19982	1.23937	1.13835
IR2	1.59035	1.41483	1.47336	1.27564
RHO=0.5				
DET	1.50012	1.38953	1.42629	1.25817
TR	1.26785	1.20541	1.22530	1.14925
IR2	1.70151	1.50492	1.56389	1.37256
RHO=0.9				
DET	1.995815	1.19457	1.15673	1.09565
TR	1.18285	1.15161	1.15774	1.10833
IR2	1.50254	1.34775	1.37285	1.25051
(c) n=60	$X_M$	$X_{HL}$	$X_{T(.25)}$	$X_{T(.10)}$
RHO=0.0				
DET	1.49876	1.39738	1.48229	1.24534
TR	1.23394	1.18524	1.21896	1.11624
IR2	1.54449	1.41140	1.48902	1.24656
RHO=0.5				
DET	1.50496	1.33902	1.37556	1.19554
TR	1.24668	1.17201	1.20011	1.10753
IR2	1.59127	1.39919	1.49082	1.24963
RHO=0.9				
DET	1.820390	1.31018	1.18986	1.18056
TR	1.20510	1.20301	1.23672	1.12477
IR2	1.63433	1.46321	1.58119	1.27445



TABLE IV  
MONTE CARLO RELATIVE EFFICIENCIES  
(Contaminated Gaussian)

(a) n=20

	$X_M$	$X_{HL}$	$X_{T(.25)}$	$X_{T(.10)}$
RH0=0.0				
DET	1.15461	1.35893	1.31252	1.39794
IR	1.06249	1.16568	1.14341	1.18242
TR2	1.10506	1.35868	1.30235	1.39830
RH0=0.5				
DET	1.09558	1.33712	1.29458	1.38399
IR	1.10032	1.16794	1.16193	1.17845
TR2	1.29283	1.37982	1.38419	1.39142
RH0=0.9				
DET	.802783	1.29001	1.21236	1.32480
IR	1.14231	1.18302	1.20082	1.19024
TR2	1.43272	1.41259	1.47405	1.42733

(b) n=40

	$X_M$	$X_{HL}$	$X_{T(.25)}$	$X_{T(.10)}$
RH0=0.0				
DET	1.06551	1.37247	1.33618	1.38673
IR	1.03031	1.16452	1.13831	1.16875
TR2	1.05777	1.34094	1.26023	1.34693
RH0=0.5				
DET	.960764	1.31239	1.21833	1.34052
IR	1.01338	1.15765	1.11721	1.16746
TR2	1.08499	1.36161	1.27141	1.38010
RH0=0.9				
DET	.657901	1.23360	1.01560	1.21047
IR	.961540	1.14275	1.07885	1.14160
TR2	.984810	1.31557	1.18676	1.31607

(c) n=60

	$X_M$	$X_{HL}$	$X_{T(.25)}$	$X_{T(.10)}$
RH0=0.0				
DET	.962990	1.37380	1.27929	1.38588
IR	.984854	1.17215	1.13370	1.17873
TR2	.976645	1.37408	1.29103	1.39275
RH0=0.5				
DET	.977959	1.31787	1.21396	1.33838
IR	1.00302	1.16438	1.11388	1.17250
TR2	1.02884	1.38654	1.26207	1.40412
RH0=0.9				
DET	.571742	1.25365	1.06703	1.34574
IR	.929473	1.16485	1.09669	1.18669
TR2	.942395	1.37158	1.22350	1.41662





TABLE X  
MONTE CARLO RELATIVE EFFICIENCIES  
(Cauchy Distribution)

(a) n=20	$X_M$	$X_{HL}$	$X_T(.25)$	$X_T(.10)$
RHO=0.0				
DET	304.650	188.950	255.633	84.0988
TR	17.8619	14.1882	16.5260	9.27750
IR2	331.226	212.282	288.761	87.6382
RHO=0.5				
DET	272.308	183.539	233.000	85.9678
TR	18.7090	14.9111	17.0922	10.7811
IR2	394.272	241.423	323.769	135.942
RHO=0.9				
DET	188.650	139.215	180.540	77.7531
TR	19.5791	15.0365	17.5623	10.7776
IR2	437.187	241.152	332.931	121.832

(b) n=40	$X_M$	$X_{HL}$	$X_T(.25)$	$X_T(.10)$
RHO=0.0				
DET	4703.87	3306.78	4469.18	1860.26
TR	74.7018	63.0659	72.4382	47.6102
IR2	6246.07	4507.03	5821.37	2600.33
RHO=0.5				
DET	4271.34	3124.83	4070.42	1673.70
TR	85.8109	72.6329	83.2610	52.3958
IR2	9396.72	6605.31	8747.66	3356.08
RHO=0.9				
DET	2824.02	2571.87	3210.32	1393.72
TR	81.3385	70.0023	79.3516	51.4988
IR2	7475.30	5244.65	6779.86	2837.74

(c) n=60	$X_M$	$X_{HL}$	$X_T(.25)$	$X_T(.10)$
RHO=0.0				
DET	3648.02	2620.04	3391.63	1676.81
TR	86.0599	71.9088	82.3380	57.6595
IR2	9672.69	6586.49	8730.48	4251.33
RHO=0.5				
DET	3487.56	2465.04	3133.91	1574.62
TR	101.144	83.6789	95.4133	66.6214
IR2	13446.5	8976.60	11872.1	5658.13
RHO=0.9				
DET	2073.92	2237.82	2650.22	1491.26
TR	88.0925	75.7093	87.1780	60.1277
IR2	8899.59	6050.33	8162.84	3789.70



TABLE XI  
MONTE CARLO RELATIVE EFFICIENCIES  
(Uniform Distribution)

(a) n=20	$X_M$	$X_{HL}$	$X_T(.25)$	$X_T(.10)$
$PHQ=0.0$				
DET	.350364	.811867	.509748	.745143
TF	.596833	.897695	.717291	.863567
TF2	.361831	.800465	.518986	.746303
$PHQ=0.5$				
DET	.369075	.844673	.568212	.788225
TF	.617849	.925703	.761551	.891987
TF2	.382659	.866939	.589724	.801619
$PHQ=0.9$				
DET	.294367	.880319	.577459	.825923
TF	.587471	.931475	.729648	.878802
TF2	.355801	.865695	.526191	.764746

(b) n=40	$X_M$	$X_{HL}$	$X_T(.25)$	$X_T(.10)$
$PHQ=0.0$				
DET	.371985	.859955	.516797	.720415
TF	.603420	.928495	.715492	.848430
TF2	.357141	.864168	.507412	.719279
$PHQ=0.5$				
DET	.412683	.831858	.562422	.751189
TF	.666271	.920458	.767215	.878550
TF2	.465439	.856487	.605448	.784533
$PHQ=0.9$				
DET	.328218	.844290	.634603	.796170
TF	.675503	.926840	.777196	.881797
TF2	.484647	.860880	.600206	.775370

(c) n=60	$X_M$	$X_{HL}$	$X_T(.25)$	$X_T(.10)$
$PHQ=0.0$				
DET	.354980	.873640	.511024	.710144
TF	.599004	.935486	.715024	.843225
TF2	.362346	.876480	.511471	.711826
$PHQ=0.5$				
DET	.386571	.842992	.546711	.751676
TF	.621918	.921413	.748817	.869483
TF2	.410950	.854102	.573252	.759658
$PHQ=0.9$				
DET	.344519	.861020	.556791	.762904
TF	.626382	.919192	.736528	.857937
TF2	.401596	.842526	.540377	.732188



## REFERENCES

- Andrews, D.F., Bickel, P.J., Hampel, F.R., Huber, P.J., Rogers, W.H., and Tukey, J.W. (1972). Robust Estimates of Location: Survey and Advances, Princeton University Press.
- Bickel, P.J. (1964). On some alternative estimates for shift in the p-variate one sample problem. Ann. Math. Statist., 35, 1079-1090.
- Bickel, P.J. (1965). On some robust estimates of location. Ann. Math. Statist., 36, 847-858.
- Gentleman, W.M. (1965). Robust estimation of multivariate location by minimizing pth power deviations. Unpublished Ph.D. thesis, Princeton University.
- Gnanadesikan, R., and Kettenring, J.R. (1972). Robust estimates, residuals, and outlier detection with multiresponse data. Biometrics, 28, 81-124.
- Hodges, J.L., and Lehmann, E.L. (1963). Estimates of location based on rank tests. Ann. Math. Stat., 34, 598-611.
- Huber, P.J. (1964). Robust estimation of a location parameter. Ann. Math. Stat., 35, 73-101.
- Huber, P.J. (1972). Robust statistics: a review. Ann. Math. Statist., 43, 1041-1067.
- Mood, A.M. (1941). On the joint distribution of the medians in samples from a multivariate population. Ann. Math. Statist., 12, 268-278.
- Tukey, J.W. (1960). A survey of sampling from contaminated distributions. In Contributions to Probability and Statistics (I. Olkin et al., eds.), Stanford University Press, 448-85.











Date Due

DEC 19 1986

1986

1986

JUL 01 1992

APR

Lib-26-67

HD28.M414 no.1133- 80  
Schmalensee, R/The new industrial orga  
739568 D\*BKS 00132000



3 9080 001 988 648

HD28.M414 no.1134- 80  
Banks, Louis. /Living in a newsocracy  
740326 D\*BKS 00137171



3 9080 002 050 042

HD28.M414 no.1135- 80  
Wong, M. Antho/A Monte Carlo study of  
740340 D\*BKS 00133429



3 9080 002 005 863

HD28.M414 no.1136- 80  
Henriksson, Ro/Tests of market timing  
740309 D\*BKS 00132385



3 9080 001 992 954

HD28.M414 no.1140- 80  
Schmalensee, R/Product differentiation  
740333 D\*BKS 00132649



3 9080 001 996 187

HD28.M414 no.1141- 80  
Buer, Terje K./Investigation of make o  
740829 D\*BKS 00132383



3 9080 001 992 913

HD28.M414 no.1142- 80  
Nevis, Edwin C/Gestalt awareness proce  
740329 D\*BKS 00132363



3 9080 001 992 558

HD28.M414 no.1143- 80  
Kolb, Deborah /Onstage with labor medi  
740812 D\*BKS 00132361



3 9080 001 992 525

HD28.M414 no.1144- 80  
Berndt, Ernst /Short run labor product  
740816 D\*BKS 00132384



3 9080 001 992 939



3 9080 004 178 924

1145-80

HD28.M414 no.1146- 80  
Wallace, Anne /Getting medical attent  
740836 D\*BKS 00132387



3 9080 001 992 996

